

Elliptic Islands on the Elliptical Stadium

Sylvie Oliffson Kamphorst Sônia Pinto de Carvalho
 Departamento de Matemática, ICEx, UFMG
 C.P. 702, 30161-970, Belo Horizonte, Brasil.

Abstract

We investigate the existence of elliptic islands for a special family of periodic orbits of a two-parameter family of maps $T_{a,h}$, corresponding to the billiard problem on the elliptical stadium.

The hyperbolic character of those orbits were studied in [2] for $1 < a < \sqrt{2}$ and here we look for the elliptical character for every $a > 1$.

We prove that, for $a < \sqrt{2}$, the lower bound for chaos $h = H(a)$ found in [2] is the upper bound of ellipticity for this special family.

For $a > \sqrt{2}$ we prove that there is no upper bound on h for the existence of elliptic islands.

The main results we use are Birkhoff Normal Form and Moser's Twist Theorem.

1 Introduction

The elliptical stadium is a plane region bounded by a curve Γ , constructed by joining two half-ellipses, with major axes $a > 1$ and minor axes $b = 1$, by two straight segments of equal length $2h$ (see fig. 1).




Figure 1: The elliptical stadium.

The billiard on the elliptical stadium consists in the study of the free motion of a point particle inside the stadium, being reflected elastically at the impacts with Γ . Since the motion is free inside Γ , it is determined either by two consecutive points of reflection at Γ or by the point of reflection and the direction of motion immediately after each collision.

For fixed a and h , let $s \in [0, L)$ be the arc length parameter for Γ and the direction of motion be given by the angle β with the normal to the boundary at the impact point. The billiard defines an invertible map $T_{a,h}$ from the annulus $\mathcal{A} = [0, L) \times (-\pi/2, \pi/2)$ into itself, preserving the measure $d\mu = \cos \beta \, d\beta \, ds$.

⁰AMS subject classification: 37E40, 70K42

Since Γ is globally C^1 but not C^2 , $T_{a,h}$ is a homeomorphism (see, for instance, [6]) and if (s_0, β_0) and $(s_1, \beta_1) = T_{a,h}(s_0, \beta_0) \in \mathcal{A}$ such that Γ is analytic in some neighborhoods of s_0 and s_1 , then clearly $T_{a,h}$ is analytic in some neighborhoods of (s_0, β_0) and (s_1, β_1) .

For each (a, h) , $(\mathcal{A}, \mu, T_{a,h})$ defines a discrete dynamical system, whose dynamics depends on the values of a and h . For instance, when $h = 0$ we have an ellipse and the billiard is integrable.

When $h \neq 0$, two main features appear. If $a < \sqrt{2}$, Donnay [3] proved that the billiard on the elliptical stadium is chaotic (in the sense of non-vanishing Lyapunov exponents) when h is sufficiently large. Lower bounds for h for this behaviour were found by Markarian and ourselves in [5] and by Canale, Markarian and ourselves in [2]. In the present work we show that the lower bound found in [2] is optimal, in the sense that below it we can assure the existence of elliptic islands of positive measure.

In [1] Bunimovich conjectured the existence of a stable periodic orbit, with island of positive measure, for billiards such as the elliptical stadium with $a > \sqrt{2}$ and $h \neq 0$. In this work we make some progress in this direction, proving that there is no upper bound on h for the existence of elliptic islands if $a > \sqrt{2}$. So, there is no way to destroy the elliptic islands by just increasing the distance between the half-ellipses.

To prove the existence of elliptic islands we extend the results about a special family of periodic orbits, called pantographic, studied in [2] and find regions on the parameter plane where at least one of its members is elliptic and stable, so, with islands of positive measure in phase space.

2 Pantographic orbits: existence and ellipticity

In this section we define the special family of periodic orbits and investigate the existence and ellipticity of its members. This family has already been investigated in [2] for $a < \sqrt{2}$. Here we extend that work for all $a > 1$. We will skip most of the proofs which can be found in the work cited above.

Given a , h and a positive integer n , an (n, a, h) -pantographic orbit, denoted by $Pan(n, a, h)$, is a symmetric $(4 + 2n)$ -periodic orbit, with exactly 2 impacts at each half-ellipse, joined by a vertical path, and crossing any vertical line only twice. One example can be seen in figure 2.




Figure 2: The 10-periodic pantographic orbit ($n = 3$).

Let the right half-ellipse of the stadium be parametrized by $(x, y) = (a \cos \lambda + h, \sin \lambda)$ and P be the point marked on figure 2. Using the obvious symmetries (see figure 3), the parameter λ of P must satisfy:

$$\tan 2\beta = \frac{a \tan \lambda}{a^2 \tan^2 \lambda - 1} = \frac{h + a \cos \lambda}{n + \sin \lambda} \quad \text{and} \quad \tan \beta = \frac{\cos \lambda}{a \sin \lambda} \quad (1)$$

where $\beta > 0$ is, as defined above, the angle of the trajectory from P , with the normal to the boundary.




Figure 3:

The following proposition gives the region of existence of those pantographic orbits in the parameter plane

Proposition 1

- $Pan(0, a, h)$ and $Pan(1, a, h)$ exist for every $a > 1$ and $h > 0$.
- For $n \geq 2$, $Pan(n, a, h)$ exists for every $1 < a \leq 2$ and $h > 0$ or for every $a > 2$ and $h > (n-1)\sqrt{a(a-2)}$.

Proof: Equation (1) can be written as

$$n = \frac{a^2 t^2 - 1}{2at} h + \frac{(a^2 - 2)t^2 - 1}{2t\sqrt{1+t^2}}. \quad (2)$$

where $t = \tan \lambda$. As proven in [2], for each integer $n \geq 0$, this equation has a unique solution $t(n, a, h) = \tan \lambda(n, a, h) > 0$ for every $a > 1$ and $h > 0$ and $t(n, a, h) \in (\frac{1}{a}, +\infty)$.

For $n = 0$ and $n = 1$ this results implies the existence of the corresponding $Pan(n, a, h)$ for every $a > 1$ and $h > 0$.

However, for $n \geq 2$ one must also ask that

$$\tan 2\beta \geq \frac{a \cos \lambda}{1 + \sin \lambda}$$

in order to guarantee that the next impact point from P is on the straight part of the boundary. This is equivalent to

$$0 \leq \sin \lambda \leq \frac{1}{a-1} \quad (3)$$

which is always true if $1 < a \leq 2$.

If $a > 2$ we rewrite (3) as $0 \leq t = \tan \lambda \leq \frac{1}{\sqrt{a(a-2)}}$. Since $\frac{\partial t}{\partial h}(n, a, h) < 0$ (which is easily verified from (2)), and $\frac{1}{a} < \frac{1}{\sqrt{a(a-2)}}$, there exists a unique $\bar{h}$ such that $t(n, a, \bar{h}) = \frac{1}{\sqrt{a(a-2)}}$ and if $h > \bar{h}$, $\frac{1}{a} < t(n, a, h) < \frac{1}{\sqrt{a(a-2)}}$. From (2), $\bar{h} = (n-1)\sqrt{a(a-2)}$. $\clubsuit$

For each fixed n , we denote by U_n the open region in the parameter plane where $Pan(n, a, h)$ exists, according to Proposition 1. For $(a, h) \in U_n$, let s be the arc length corresponding to the point P of $Pan(n, a, h)$ and β the angle with the normal of the trajectory at this point as before. Then $T_{a,h}^{4+2n}(s, \beta) = (s, \beta)$ and the ellipticity of this orbit is determined by the eigenvalues of $\mathcal{D}T_{a,h}^{4+2n}|_{(s,\beta)}$.

As shown in [2], we can write $\mathcal{D}T_{a,h}^{4+2n}|_{(s,\beta)} = (M_1 M_2)^2$ with

$$M_j = \frac{1}{\cos \beta} \begin{pmatrix} l_j K - \cos \beta & l_j \\ K(l_j K - 2 \cos \beta) & l_j K - \cos \beta \end{pmatrix}$$

and where l_1 is the length of the trajectory between two impacts with the same half-ellipse, l_2 is the length of the trajectory between two impacts with the different half-ellipses and K is the curvature of the ellipse at s .

Let

$$\Delta_n(a, h) = \left(\frac{l_1 K}{\cos \beta} - 1 \right) \left(\frac{l_2 K}{\cos \beta} - 1 \right).$$

Since $\det(M_1 M_2) = 1$ and $\text{tr}(M_1 M_2) = 4\Delta_n(a, h) - 2$, it follows that if $0 < \Delta_n(a, h) < \frac{1}{2} < \Delta_n(a, h) < 1$ then $Pan(n, a, h)$ is *elliptic* (meaning that the eigenvalues of $\mathcal{D}T_{a,h}^{4+2n}|_{(s,\beta)}$ are unitary with non zero imaginary part).

The following lemma summarizes some properties of $\Delta_n(a, h)$ and its technical proof has been postponed to the appendix.

Lemma 2 *For every $n \geq 0$, let*

$$\tilde{U}_n = \{(a, h) | 1 < a < \sqrt{2}, h > 0\} \cup \{(a, h) | a \geq \sqrt{2}, h > na\sqrt{a^2 - 2} \equiv h_n^0(a)\} \subset U_n.$$

The function $\Delta_n(a, h)$ has the following properties:

1. $\Delta_n(a, h)|_{\tilde{U}_n} > 0$
2. $\frac{\partial \Delta_n}{\partial h}|_{\tilde{U}_n} > 0$
3. $\lim_{h \rightarrow +\infty} \Delta_n(a, h) = +\infty$
4. for $1 < a < \sqrt{2}$, $\lim_{h \rightarrow 0} \Delta_n(a, h) = L_n(a) = (\frac{2}{a^2} - 1)(\frac{2(n+1)}{a^2} - 1) > 0$
for $a \geq \sqrt{2}$, $\lim_{h \rightarrow na\sqrt{a^2 - 2}} \Delta_n(a, h) = 0$.

For each $0 < c \leq 1 + 2n$, let α_n^c be the unique solution of $L_n(a) = c$. We have that $1 < \alpha_n^c < \sqrt{2}$ and if $c < d$ then $\alpha_n^d < \alpha_n^c$.

It follows from the lemma that every level curve $\Delta_n = c$ is given by a graph $h_n^c : (\alpha_n^c, +\infty) \rightarrow \mathbb{R}$ such that $\Delta_n(a, h_n^c(a)) = c$ and $\Delta_n(a, h) < c$ if $h < h_n^c(a)$ and $\Delta_n(a, h) > c$ if $h > h_n^c(a)$.

The characterization of the region of ellipticity is then given by:




Figure 4: Region of ellipticity with lines of resonance up to order 4 for $n = 3$

Proposition 3 *For each fixed n , the region in the parameter plane where $Pan(n, a, h)$ is elliptic is the union of two open adjacent strips, bounded by the graphs of $h_n^0(a), h_n^{1/2}(a), h_n^1(a)$ and by the segments $\{(a, 0) | \alpha_n^1 < a < \alpha_n^{1/2}\}$ and $\{(a, 0) | \alpha_n^{1/2} < a < \sqrt{2}\}$. (see figure 4).*

For each n and (a, h) in the region of ellipticity of $Pan(n, a, h)$, let μ and $\bar{\mu}$ be the eigenvalues of $DT_{a,h}^{4+2n}|_{(s,\beta)}$. $Pan(n, a, h)$ has a resonance of order k if $\mu^k = 1$, i.e., $\mu = e^{i\frac{2j\pi}{k}}, j = 1, \dots, k-1$. It is easy to show that this corresponds, for $k > 1$, to

$$\Delta_n(a, h) = \frac{1}{2} \left(1 + \cos \frac{j\pi}{k} \right) \equiv c_{jk}.$$

Clearly $0 < c_{jk} < 1$ and we have the curves of resonance given by the graphs of $h_n^{c_{jk}} : (\alpha_n^{c_{jk}}, +\infty) \rightarrow \mathbb{R}$.

We can then characterize the region of ellipticity with no resonances up to order k by:

Proposition 4 *For a fixed n , the region in the parameter plane where $Pan(n, a, h)$ is elliptic with no resonances up to order k is a finite union of open disjoint adjacent strips contained in the region of ellipticity. (see figure 4)*

3 Existence of elliptic islands

In order to establish if the elliptic periodic orbits described in the previous section have invariant curves surrounding them, we will invoke the two classical results:

Birkhoff Normal Form: *Let f be an area preserving map in C^l ($l \geq 4$) with a fixed point at the origin, with eigenvalues μ and $\bar{\mu}$, $|\mu| = 1$. If for some integer q in $4 \leq q \leq l+1$ one has $\mu^k \neq 1$ for $k = 1, 2, \dots, q$ then there exists a real analytic transformation taking f into the normal form*

$$\zeta \rightarrow f(\zeta, \bar{\zeta}) = \mu \zeta e^{i\tau(\zeta, \bar{\zeta})} + g(\zeta, \bar{\zeta}) \quad (4)$$

where $\tau(\zeta\bar{\zeta}) = \tau_1|\zeta|^2 + \dots + \tau_s|\zeta|^{2s}$, with $s = [\frac{q}{2}] - 1$, is a real polynomial in $|\zeta|^2$ and g vanishes with its derivatives up to order $q - 1$ at $\zeta = \bar{\zeta} = 0$.

Theorem (Moser, [8], p.56) *If the polynomial $\tau(|\zeta|^2)$ does not vanishes identically, $\zeta = 0$ is a stable fixed point (which means that there are invariant curves surrounding it, and so an elliptic island of positive measure).*

For each fixed period n_0 , we will then investigate the resonances of $T_{a,h}^{4+2n_0}$ and the zeros of the coefficients of its Birkoff normal form, near the Pantographic orbit.

Let us fix a period n_0 and a major axis $a_0 > \alpha_{n_0}^1 = \sqrt{(2+2n_0)/(2+n_0)}$. According to proposition 4, $Pan(n_0, a_0, h)$ is elliptic with no resonances up to order q if h is in the finite union of open disjoint adjacent intervals denoted $\cup I_j^q$.

Let $\lambda(h), s(h), \beta(h)$ be respectively the parameter, the arc length and the angle with the normal for the point P of $Pan(n_0, a_0, h)$.

Lemma 5 *For any $q \geq 4$, $\lambda(h), s(h)$ and $\beta(h)$ are analytic functions of h on each I_j^q .*

Proof: The point P will belong to $Pan(n_0, a_0, h)$ if $t = \tan \lambda$ satisfies equation (2):

$$n_0 = \frac{a_0^2 t^2 - 1}{2a_0 t} h + \frac{(a_0^2 - 2)t^2 - 1}{2t\sqrt{1+t^2}}.$$

$A(t) = \frac{a_0^2 t^2 - 1}{2a_0 t}$ and $B(t) = \frac{(a_0^2 - 2)t^2 - 1}{2t\sqrt{1+t^2}}$ are analytical functions of t and $A(t) \neq 0$, since $\frac{1}{a_0} < t < \infty$. So $h = h(t) = \frac{n_0 - B(t)}{A(t)}$ is analytic. As this equation has a unique solution for each h , the inverse $t = t(h)$ exists and is then locally analytic for every $h \in I_j^q$.

The functions $\lambda(h) = \arctan t(h)$ and the corresponding arc length of the ellipse $s(h) = s(\lambda(h))$ are then analytic. The same is true for $\beta(h) = \beta(\lambda(h))$. ♣

For each fixed h , let f be the translation of $T_{a_0,h}^{4+2n_0}$ by $(s(h), \beta(h))$. The map f is clearly area preserving and analytic in (s, β) on a neighbourhood of the origin. The eigenvalues of $Df_{(0,0)}$ are the same as those of $DT_{a_0,h}^{4+2n_0}(s(h), \beta(h))$. If $h \in I_j^q$, f can be written in the Birkhoff normal form (4). If one of the Birkhoff coefficients is not zero, f has an elliptic island surrounding $(0, 0)$ and, by translation, there is an elliptic island surrounding $Pan(n_0, a_0, h)$.

Lemma 6 *For $1 \leq m \leq [\frac{q}{2}] - 1$, $\tau_m(h)$, the m -th Birkhoff coefficient of f , is an analytic function of h on each I_j^q .*

Proof: For each fixed h , $\tau_m(h)$ is a combination of the coefficients of the $(q - 1)$ -th jet of f at $(0, 0)$ (see, for instance, [7]). The steps giving rise to the calculation of $\tau_m(h)$ (complexification of the space, elimination of unwanted terms) are analytical. So, if the coefficients of the jet $J_{q-1}f_{(0,0)}$ are analytical in h , then $\tau_m(h)$ will also be an analytic function of h .

Now these coefficients are combinations of the entries of $DT_{a_0,h}^{4+2n_0}$ and their derivatives up to $(q - 1)$ -th order with respect to s and β , calculated at $(s(h), \beta(h))$.

Let (s, β) and (s', β') be two consecutive impacts of a trajectory with the two different half-ellipses (with $l \geq 0$ impacts with the straight parts between them), or two consecutive impacts of a trajectory with the

same half-ellipse (with $l = 0$). Then $DT_{a_0, h}^{4+2n_0}(s, \beta)$, for (s, β) near to $(s(h), \beta(h))$ is a finite product of matrices of the form (see, for instance, [6])

$$\frac{(-1)^l}{\cos \beta'} \begin{pmatrix} LK - \cos \beta & L \\ KK'L - K' \cos \beta - K \cos \beta' & LK' - \cos \beta' \end{pmatrix}$$

where K stands for the curvature of the ellipse at the impact and L is the total length of the trajectory between the two impacts with the half-ellipses.

Since $\cos \beta \neq 0$ for β near $\beta(h)$, all the entries of the matrix above, as well as their derivatives up to any order in s and β , are analytic functions of h . Using lemma 5 we conclude that all the coefficients of $J_{q-1}f_{(0,0)}$ are analytic in h .

It follows that $\tau_m(h)$ is analytic in h , leading immediately to the next corollary. ♣

Corollary 7 *On each I_j^q and for $1 \leq m \leq [\frac{q}{2}] - 1$, the set $\{h / \tau_m(h) = 0\}$ is either the entire I_j^q or a discrete set.*

In order to prove the existence of islands we use the natural recurrence on the order of the resonances.

We begin by analysing the zeros of τ_1 on the non resonant intervals I_j^4 . If $\tau_1 = 0$ only on a discrete subset of each I_j^4 , $Pan(n_0, a_0, h)$ has elliptic island except for a discrete set of values of h (which can be smaller than the union of the discrete subsets of zeros of τ_1 and the values of resonance up to order 4, since on the discrete subsets a non resonant value of higher order may have a non zero Birkhoff coefficient).

If τ_1 is identically zero on one of those intervals, we proceed to the next step, applying the same analysis to the zeros of τ_2 .

We continue the recurrence and it will end up in a finite number of steps if for some order of resonance the last Birkhoff coefficient does not vanish identically on a whole non resonant interval. Otherwise, all the Birkhoff coefficients will vanish on at least one open interval, bounded by resonant values of h . In this last case, since $\frac{\partial \Delta_n}{\partial h} > 0$, the rotation number $\rho(h)$ of $Pan(n_0, a_0, h)$ is not constant and there exists h_0 such that $\rho(h_0)$ is diophantine. As f is analytic, it is conjugate to a rotation and there will be invariant curves [4].

We conclude that:

Theorem 1 *Given n and $a > \alpha_n^1$, there are at least countably many values of h in $\cup I_j^4$ such that $Pan(n, a, h)$ has an elliptic island.*

Remark: As Moeckel proved in [7], in a generic one-parameter family of area preserving maps with elliptic fixed points, the first Birkhoff coefficient τ_1 varies from $-\infty$ to $+\infty$ as the rotation number varies from 0 to 1/3 or from 2/3 to 1. So, we do not expect $\tau_1(h)$ to be always different from zero, neither to vanish identically. Furthermore, generically, at a zero of τ_1 , a higher Birkhoff coefficient will not vanish. So, although we can handle the case $\tau_s(h_0) = 0, \forall s$, we do not expect it to happen in our case.

4 Bounds for the existence of islands

4.1 The case $a < \sqrt{2}$

As shown in proposition 3 and in [2], if (a, h) is in the region of ellipticity of $Pan(n, a, h)$ then $a \geq \alpha_n^1 = \sqrt{\frac{2n+2}{n+2}}$. So, given $a \in (1, \sqrt{2})$, there is only a finite number of periods $4 + 2n$ such that $Pan(n, a, h)$ can be elliptic. More precisely, $n \leq \frac{2(a^2-1)}{2-a^2}$.

Let $H(a)$ be the maximum of $h_n^1(a)$ for those periods. As proved in [2], $H(a)$ is a lower bound for chaos. By theorem 1, it is also an upper bound for the existence of elliptic islands for the Pantographic family.

4.2 The case $a > \sqrt{2}$

On the other hand, if $a > \sqrt{2}$, $Pan(n, a, h)$ can be elliptic for any period n . Moreover, for each n and q , $\cup I_j^q \subset (h_n^0(a), h_n^1(a))$, with $h_n^0(a) = na\sqrt{a^2-2}$. We also have that $(h_n^0(a), h_n^1(a)) \cap (h_n^0(a), h_{n+1}^0(a))$ is a non empty open interval. So we can find h , $na\sqrt{a^2-2} < h < (n+1)a\sqrt{a^2-2}$ such that $Pan(n, a, h)$ has an elliptic island. This proves the following

Theorem 2 *Given $a > \sqrt{2}$ there is no upper bound on h for the existence of elliptic islands on the elliptical stadium billiard.*

However, as can be seen in figure 5, for values of a further away from $\sqrt{2}$, the strips of ellipticity are disjoint. In these gaps all pantographic orbits are hyperbolic, having, thus, no islands.




Figure 5: Gaps between the strips of ellipticity

Nevertheless, in our simulations other islands appear, obviously corresponding to different periodic orbits. In figure 6 we exemplify this fact for $a = 2$ and $h = 2$, a value located in the gap between the strips of ellipticity for $n = 0$ and $n = 1$ (see figure 5). We show three non-pantographic orbits and their islands and the whole phase space, where we can see many other islands surrounded by what seems to be a chaotic sea. As far as our results indicate and our simulations show, this should be the typical picture for the phase space when $a > \sqrt{2}$.




Figure 6: Phase space for $(a, h) = (2, 2)$

5 Appendix

5.1 Some properties of $t(n, a, h)$

We called U_n the open region in the parameter plane where $Pan(n, a, h)$ exists:

$$U_0 = U_1 = \{(a, h) / a > 1, h > 0\}$$

$$U_n = \{(a, h) / 1 < a \leq 2, h > 0\} \cup \{(a, h) / a > 2, h > (n-1)\sqrt{a(a-2)}\},$$

for $n \geq 2$.

The following lemma gives some useful information about $t(n, a, h)$, the solution of (1) or (2).

Lemma 8 *Given $n \geq 0$, let $(a, h) \in U_n$ and $t(n, a, h)$ be the unique solution of (2). Then*

- $\forall n$, when $h \rightarrow +\infty$, $t(n, a, h) \rightarrow \frac{1}{a}$.
- For $n = 0$
 - if $1 < a \leq \sqrt{2}$, when $h \rightarrow 0^+$, $t(0, a, h) \rightarrow +\infty$
 - if $a > \sqrt{2}$, when $h \rightarrow 0^+$, $t(0, a, h) \rightarrow \frac{1}{\sqrt{a^2-2}}$
- For $n = 1$
 - if $1 < a \leq 2$, when $h \rightarrow 0^+$, $t(1, a, h) \rightarrow +\infty$
 - if $a > 2$, when $h \rightarrow 0^+$, $t(1, a, h) \rightarrow \frac{1}{\sqrt{a(a-2)}}$
- For $n \geq 2$
 - if $1 < a \leq 2$, when $h \rightarrow 0^+$, $t(n, a, h) \rightarrow +\infty$
 - if $a > 2$, when $h \rightarrow (n-1)\sqrt{a(a-2)}^+$, $t(n, a, h) \rightarrow \frac{1}{\sqrt{a(a-2)}}$

Proof: Since $\frac{\partial t}{\partial h} < 0$ and $\frac{1}{a} < t$, $\lim_{h \rightarrow +\infty} t$ exists. From equation (1),

$$at - \frac{1}{at} = 2 \frac{n\sqrt{1+t^2} + t}{h\sqrt{1+t^2} + a} \rightarrow 0 \quad (5)$$

as $h \rightarrow +\infty$ and so $t \rightarrow \frac{1}{a}$.

To study the limit as $h \rightarrow 0^+$, let us take $x = \frac{1}{t}$. Equation (1) becomes

$$x(2n\sqrt{1+x^2} + x^2 - (a^2 - 2)) = (a^2 - x^2)\sqrt{1+x^2} \frac{h}{a}.$$

Since $0 < x < \frac{1}{a}$, $x(2n\sqrt{1+x^2} + x^2 - (a^2 - 2)) > 0$.

Let $\bar{x} = \lim_{h \rightarrow 0^+} x$. We have that $0 \leq \bar{x} \leq a$ and $\bar{x}(2n\sqrt{1+\bar{x}^2} + \bar{x}^2 - (a^2 - 2)) = 0$.

For $n = 0$, if $1 < a \leq \sqrt{2}$, $\bar{x} = 0$ is the unique solution of this equation and $\lim_{h \rightarrow 0^+} t(0, a, h) = +\infty$. If $a > \sqrt{2}$ we have a new solution $\bar{x} = \sqrt{a^2 - 2}$. But for $0 < x < \sqrt{a^2 - 2}$, $x(x^2 - (a^2 - 2)) < 0$. So $\lim_{h \rightarrow 0^+} x = \sqrt{a^2 - 2}$ and $\lim_{h \rightarrow 0^+} t(0, a, h) = \frac{1}{\sqrt{a^2 - 2}}$.

For $n = 1$ we have $\bar{x}(2\sqrt{1+\bar{x}^2} + \bar{x}^2 - (a^2 - 2)) = 0$. If $a^2 - 2 \leq 2$, i.e. $a \leq 2$, the unique solution is $\bar{x} = 0$ and $\lim_{h \rightarrow 0^+} t(1, a, h) = +\infty$. If $a > 2$ the second solution is $\bar{x} = \sqrt{a(a-2)}$. As above, if $0 < x < \sqrt{a(a-2)}$, $x(2\sqrt{1+x^2} + x^2 - (a^2 - 2)) < 0$ and so $\lim_{h \rightarrow 0^+} t(1, a, h) = \frac{1}{\sqrt{a(a-2)}}$.

For $n \geq 2$, we remark first that if $k > l$ then $t(k, a, h) > t(l, a, h)$. So, for $1 < a \leq 2$ $\lim_{h \rightarrow 0^+} t(n, a, h) = +\infty$.

For $a > 2$, the limit as $h \rightarrow (n-1)\sqrt{a(a-2)}$ is the unique solution of equation (1) for $h = (n-1)\sqrt{a(a-2)}$ which is $t = \frac{1}{\sqrt{a(a-2)}}$. ♣

Remark: When $h \rightarrow 0^+$, the elliptical stadium becomes an ellipse. $t \rightarrow +\infty$ means that the pantographic orbit goes to the elliptic periodic orbit which corresponds to the minor axis of the ellipse.

Let us call *pantographic-like* orbits in the elliptical billiard the periodic trajectories that have vertical segments both at left and right extremes. As can be seen in [5], the 4-periodic pantographic-like orbit exists if $a > \sqrt{2}$ and the 6-periodic if $a > 2$, they are parabolic and their position is given, respectively, by $t = \frac{1}{\sqrt{a^2 - 2}}$ and $t = \frac{1}{\sqrt{a(a-2)}}$. They are the calculated limits of the 4 and 6-periodic pantographic orbits of the elliptical stadium.

5.2 Proof of the lemma 2

For a fixed n , let $(a, h) \in U_n$ and $\lambda_n(a, h)$ be the solution of (1), β be the angle, with the normal, of the outgoing trajectory at $P = (a \cos \lambda_n + h, \sin \lambda_n)$ and s the corresponding arc length.

Let

$$\Delta_n(a, h) = \left(\frac{l_1 K}{\cos \beta} - 1 \right) \left(\frac{l_2 K}{\cos \beta} - 1 \right)$$

where $l_1 = 2 \sin \lambda_n$, $l_2 = 2\sqrt{(h + a \cos \lambda_n)^2 + (n + \sin \lambda_n)^2}$ and $K = a/(a^2 \sin^2 \lambda_n + \cos^2 \lambda_n)^{3/2}$.

Let $\delta_1(a, h) = \left(\frac{l_1 K}{\cos \beta} - 1 \right)$ and $\delta_2(a, h) = \left(\frac{l_2 K}{\cos \beta} - 1 \right)$.

Lemma 9 For every $n \geq 0$ the function $\delta_1(a, h)$ has the following properties:

1. $\delta_1(a, h) > 0$ for $1 < a < \sqrt{2}$ and $h > 0$.
2. $\lim_{h \rightarrow 0^+} \delta_1(a, h) = \frac{2}{a^2} - 1$ for $1 < a < \sqrt{2}$.
3. $\delta_1(a, na\sqrt{a^2 - 2}) = 0$ for $a \geq \sqrt{2}$.
4. $\frac{\partial \delta_1}{\partial h} > 0$ for $(a, h) \in U_n$.

Proof: We have $\delta_1(a, h) = \frac{l_1 K}{\cos \beta} - 1 = 2 \frac{1+t^2}{1+a^2 t^2} - 1$ and properties 1 and 2 follow immediately.

If $a \geq \sqrt{2}$, $\delta_1 = 0$ implies $(a^2 - 2)t^2 - 1 = 0$ and $t = \frac{1}{\sqrt{a^2 - 2}}$. From equation (1) $h = na\sqrt{a^2 - 2}$ and property 3 follows.

Since $\frac{\partial \delta_1}{\partial h} = \frac{\partial \delta_1}{\partial t} \frac{\partial t}{\partial h}$, as $\frac{\partial \delta_1}{\partial t} = \frac{4t(1-a^2)}{(1+a^2 t^2)^2} < 0$, for $a > 1$, and $\frac{\partial t}{\partial h} < 0$, $\frac{\partial \delta_1}{\partial h} > 0$. ♣

Lemma 10 For every $n \geq 0$ the function $\delta_2(a, h)$ has the following properties:

1. $\delta_2(a, h) > \delta_1(a, h) > 0$ for $(a, h) \in U_n$.
2. $\lim_{h \rightarrow 0^+} \delta_2(a, h) = 2 \frac{n+1}{a^2} - 1$ for $1 < a < \sqrt{2}$.
3. $\lim_{h \rightarrow +\infty} \delta_2(a, h) = +\infty$ for $1 < a$.
4. $\frac{\partial \delta_2}{\partial h} > 0$ for $(a, h) \in U_n$.

Proof: Since $l_2 > l_1$, $\delta_2 > \delta_1$.

For $1 < a < \sqrt{2}$, when $h \rightarrow 0$, $t(n, a, h) \rightarrow +\infty$. So $l_2 \rightarrow 2(n+1)$, $K \rightarrow 1/a^2$ and $\cos \beta \rightarrow 1$ and property 2 follows.

Property 3 is obvious since $l_2 \rightarrow \infty$ as $h \rightarrow \infty$ and all the other quantities are bounded.

By definition $\delta_2(a, h) = l_2(a, h, t(a, h)) \frac{K(t(a, h))}{\cos \beta(t(a, h))} - 1$ and $\frac{\partial \delta_2}{\partial h} = \frac{K}{\cos \beta} \frac{\partial l_2}{\partial h} + \frac{\partial}{\partial t} \left(\frac{l_2 K}{\cos \beta} \right) \frac{\partial t}{\partial h}$.

We have that $\frac{\partial t}{\partial h} < 0$ and $\frac{\partial t}{\partial \lambda} > 0$. The curvature $K > 0$ and for $0 < \lambda < \pi/2$, $\frac{\partial K}{\partial \lambda} < 0$. So $\frac{\partial K}{\partial t} < 0$. As $0 < \beta < \pi/4$, $\cos \beta > 0$. As $\tan \beta = 1/at$, $\frac{\partial}{\partial t} \cos \beta > 0$. This implies that $\frac{\partial}{\partial t} \left(\frac{K}{\cos \beta} \right) < 0$.

We have that $\frac{1}{4} l_2^2 = (h + a \cos \lambda)^2 + (n + \sin \lambda)^2$. So $\frac{1}{2} l_2 \frac{\partial l_2}{\partial h} = 2(h + a \cos \lambda) > 0$ in U_n , implying that $\frac{\partial l_2}{\partial h} > 0$. We also have that $\frac{1}{4} \frac{\partial l_2^2}{\partial \lambda} = -2a \sin \lambda (n + \sin \lambda) (\tan 2\beta - \tan \beta) < 0$ as $0 < \beta < \pi/4$. So $\frac{\partial l_2}{\partial \lambda} = \frac{\partial l_2}{\partial t} \frac{\partial t}{\partial \lambda} < 0$, and $\frac{\partial l_2}{\partial t} < 0$.

This shows that $\frac{l_2 K}{\cos \beta}$ is the product of two positive decreasing functions of t and so $\frac{\partial}{\partial t} \left(\frac{l_2 K}{\cos \beta} \right) < 0$.

We conclude that $\frac{\partial \delta_2}{\partial h} > 0$. ♣

We have defined $\tilde{U}_n = \{(a, h) / 1 < a < \sqrt{2}, h > 0\} \cup \{(a, h) / a \geq \sqrt{2}, h > na\sqrt{a^2 - 2}\} \subset U_n$.

Lemma 2 For every $n \geq 0$ the function $\Delta_n(a, h)$ has the following properties:

1. $\Delta_n(a, h)|_{\tilde{U}_n} > 0$
2. $\frac{\partial \Delta_n}{\partial h}|_{\tilde{U}_n} > 0$
3. $\lim_{h \rightarrow +\infty} \Delta_n(a, h) = +\infty$
4. for $1 < a < \sqrt{2}$, $\lim_{h \rightarrow 0} \Delta_n(a, h) = L_n(a) = (\frac{2}{a^2} - 1)(\frac{2(n+1)}{a^2} - 1) > 0$
for $a \geq \sqrt{2}$, $\lim_{h \rightarrow na\sqrt{a^2-2}} \Delta_n(a, h) = 0$.

Proof: In $\tilde{U}_n$, Δ_n is the product of two positive increasing functions of h and properties 1 and 2 follows. Properties 3 and 4 follow immediately from lemmas 9 and 10. ♣

Acknowledgments

We want to thank A.Chenciner, M.Herman, C.Grotta Ragazzo and R.Roussarie. We also thank the support of CNPq during this work.

References

- [1] L. A. Bunimovich: Conditions on stochasticity of two-dimensional billiards. *Chaos* **1**, 187-193, (1991)
- [2] E. Canale, R. Markarian, S. Oliffson Kamphorst, S. Pinto de Carvalho: A lower bound for chaos on the elliptical stadium. *Physica D* **115**, 189-202 (1998)
- [3] V. J. Donnay: Using integrability to produce chaos: billiards with positive entropy. *Comm. Math. Phys.* **141**, 225-257 (1991)
- [4] M. Herman: private communication
- [5] R. Markarian, S. Oliffson Kamphorst, S. Pinto de Carvalho: Chaotic Properties of the Elliptical Stadium. *Comm. Math. Phys.* **174**, 661-679 (1996)
- [6] R. Markarian: Introduction to the ergodic theory of plane billiards. In: *Dynamical Systems*, Santiago de Chile, 1990.
- [7] R. Moeckel: Generic bifurcations of the twist coefficient, *Erg.Th.Dyn.Syst.* **10**, 185-195 (1990)
- [8] J. Moser: *Stable and random motions in dynamical systems*, PUP, Princeton, 1973.

email: syok@mat.ufmg.br and sonia@mat.ufmg.br